


BRIEF COMMUNICATION

Google Trends terms reporting rhinitis and related topics differ in European countries

J. Bousquet^{1,2,3}, I. Agache⁴, J. M. Anto^{5,6,7}, K. C. Bergmann^{8,9}, C. Bachert¹⁰ , I. Annesi-Maesano^{11,12}, P. J. Bousquet^{11,12}, G. D'Amato¹³, P. Demoly¹⁴, G. De Vries¹⁵, E. Eller¹⁶, W.J. Fokkens¹⁷, J. Fonseca^{18,19}, T. Haahtela²⁰, P. W. Hellings²¹, J. Just^{22,23}, T. Keil^{24,25}, L. Klimek²⁶, P. Kuna²⁷, K. C. Lodrup Carlsen^{28,29}, R. Mösges³⁰, R. Murray³¹, K. Nekam³², G. Onorato¹, N. G. Papadopoulos^{33,34}, B. Samolinski³⁵, P. Schmid-Grendelmeier³⁶, M. Thibaudon³⁷, P. Tomazic³⁸, M. Triggiani³⁹, A. Valiulis^{40,41}, E. Valovirta^{42,43}, M. Van Eerd¹⁵, M. Wickman^{44,45}, T. Zuberbier^{8,9} & A. Sheikh⁴⁶

¹MACVIA-France, Contre les MALadies Chroniques pour un Vieillessement Actif en France European Innovation Partnership on Active and Healthy Ageing Reference Site, Montpellier; ²INSERM U 1168, VIMA: Ageing and chronic diseases Epidemiological and public health approaches, Villejuif; ³UMR-S 1168, Université Versailles St-Quentin-en-Yvelines, Montigny le Bretonneux, France; ⁴Faculty of Medicine, Transylvania University, Brasov, Romania; ⁵ISGLoBAL, Centre for Research in Environmental Epidemiology (CREAL); ⁶IMIM (Hospital del Mar Research Institute); ⁷CIBER Epidemiología y Salud Pública (CIBERESP), Universitat Pompeu Fabra (UPF), Barcelona, Spain; ⁸Department of Dermatology and Allergy, Comprehensive Allergy-Centre-Charité, Charité – Universitätsmedizin Berlin; ⁹Global Allergy and Asthma European Network (GA²LEN), Berlin, Germany; ¹⁰Upper Airways Research Laboratory, ENT Department, Ghent University Hospital, Ghent, Belgium; ¹¹EPAR U707 INSERM; ¹²EPAR UMR-S UPMC, Paris VI, Paris, France; ¹³Division of Respiratory and Allergic Diseases, Hospital 'A Cardarelli', University of Naples Federico II, Naples, Italy; ¹⁴Department of Respiratory Diseases, Montpellier University Hospital, Montpellier, France; ¹⁵Peercode DV, Amsterdam, The Netherlands; ¹⁶Department of Dermatology and Allergy Centre, Odense University Hospital, Odense, Denmark; ¹⁷Department of Otorhinolaryngology, Academic Medical Centre, Amsterdam, The Netherlands; ¹⁸Center for Health Technology and Services Research- CINTESIS, Faculdade de Medicina, Universidade do Porto; ¹⁹Allergy Unit, CUF Porto Instituto & Hospital, Porto, Portugal; ²⁰Skin and Allergy Hospital, Helsinki University Hospital, Helsinki, Finland; ²¹Laboratory of Clinical Immunology, Department of Microbiology and Immunology, KU Leuven, Leuven, Belgium; ²²Allergology Department, Centre de l'Asthme et des Allergies Hôpital d'Enfants Armand-Trousseau (APHP); ²³Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Equipe EPAR, Paris, France; ²⁴Institute of Social Medicine, Epidemiology and Health Economics, Charité – Universitätsmedizin Berlin, Berlin; ²⁵Institute for Clinical Epidemiology and Biometry, University of Wuerzburg, Wuerzburg; ²⁶Center for Rhinology and Allergology, Wiesbaden, Germany; ²⁷Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Lodz, Poland; ²⁸Department of Paediatrics, Oslo University Hospital; ²⁹Faculty of Medicine, Institute of Clinical Medicine, University of Oslo, Oslo, Norway; ³⁰Institute of Medical Statistics, Informatics and Epidemiology, Medical Faculty, University of Cologne, Cologne, Germany; ³¹Medical Communications Consultant, MedScript Ltd, Dundalk, Co Louth, Ireland; ³²Hospital of the Hospitaller Brothers in Buda, Budapest, Hungary; ³³Center for Pediatrics and Child Health, Institute of Human Development, Royal Manchester Children's Hospital, University of Manchester, Manchester, UK; ³⁴Allergy Department, 2nd Pediatric Clinic, Athens General Children's Hospital 'P&A Kyriakou', University of Athens, Athens, Greece; ³⁵Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Warsaw, Poland; ³⁶Allergy Unit, Department of Dermatology, University Hospital of Zurich, Zurich, Switzerland; ³⁷RNSA (Réseau National de Surveillance Aérobiologique), Brussieu, France; ³⁸Department of ENT, Medical University of Graz, Graz, Austria; ³⁹Division of Allergy and Clinical Immunology, University of Salerno, Salerno, Italy; ⁴⁰Vilnius University Clinic of Children's Diseases and Public Health Institute, Vilnius, Lithuania; ⁴¹European Academy of Paediatrics (EAP/UEMS-SP), Brussels, Belgium; ⁴²Department of Lung Diseases and Clinical Allergology, University of Turku; ⁴³Allergy Clinic, Terveystalo, Turku, Finland; ⁴⁴Sachs' Children and Youth Hospital, Södersjukhuset; ⁴⁵Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; ⁴⁶Centre of Medical Informatics, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK

To cite this article: Bousquet J, Agache I, Anto JM, Bergmann KC, Bachert C, Annesi-Maesano I, Bousquet PJ, D'Amato G, Demoly P, De Vries G, Eller E, Fokkens WJ, Fonseca J, Haahtela T, Hellings PW, Just J, Keil T, Klimek L, Kuna P, Lodrup Carlsen KC, Mösges R, Murray R, Nekam K, Onorato G, Papadopoulos NG, Samolinski B, Schmid-Grendelmeier P, Thibaudon M, Tomazic P, Triggiani M, Valiulis A, Valovirta E, Van Eerd M, Wickman M, Zuberbier T, Sheikh A. Google Trends terms reporting rhinitis and related topics differ in European countries. *Allergy* 2017; **72**: 1261–1266.

Keywords

allergy; asthma; Google Trends; hay fever; rhinitis.

Abstract

Google Trends (GT) searches trends of specific queries in Google and reflects the real-life epidemiology of allergic rhinitis. We compared Google Trends terms related to allergy and rhinitis in all European Union countries, Norway and Switzerland from 1 January 2011 to 20 December 2016. The aim was to assess

Correspondence

Professor Jean Bousquet, CHRU Arnaud de Villeneuve, 371 Avenue du Doyen Gaston Giraud, 34295 Montpellier Cedex 5, France. Tel.: +33 611 42 88 47
E-mail: jean.bousquet@orange.fr

Accepted for publication 25 January 2017

DOI:10.1111/all.13137

Edited by: Hans-Uwe Simon

Forecasting the onset and duration of the pollen season is of vital importance in the management of allergic rhinitis (1, 2). Pollen counts may assess pollen exposure (1–3), but do not necessarily correlate with individual patients. Allergen content in the air can be assessed (4), but this requires sophisticated methods. Meteorological data may be of interest, but more research is needed (5). Combining several sources using advanced data engineering may also be important, but this is not yet available for all pollen species (6).

As forecasting is still a limited option, real-time epidemiological surveillance is an alternative option. Google Trends (GT), a Web-based surveillance tool, uses Google to explore the searching trends of specific queries. It can assess the seasonality of allergic diseases (7–10) and reflects the real-world epidemiology of symptomatic allergic rhinitis (11). However, cultural differences exist between countries, and the same search terms may not be used across all countries, making comparisons difficult. MASK (MACVIA-ARIA Sentinel NetworkK) (12) proposed to develop a pan-European sentinel network to inform allergy sufferers of the onset of the pollen season (1), and GT may be one of the methods used.

This study was conceived as a preliminary step to explore the use of GT in the epidemiological surveillance of AR and was aimed at assessing whether there are differences in the Google searches of AR and related topics. We analysed multiple time series of GT search data on rhinitis, allergy and asthma in the 28 European Union (EU) countries, Norway and Switzerland from 2011 to 2016. The aims were (i) to assess whether GT could report rhinitis and related topics accurately in all the countries; (ii) to assess the seasonality of queries; and (iii) to determine whether terms differed between countries.

Methods

The following terms were used: 'rhinitis', 'allergic rhinitis', 'hay fever', 'asthma', 'pollen', 'allergy', 'conjunctivitis' (disease, topic and term). However, only the following five

whether the same terms could be used to report the seasonal variations of allergic diseases. Using the Google Trend 5-year graph, an annual and clear seasonality of queries was found in all countries apart from Cyprus, Estonia, Latvia, Lithuania and Malta. Different terms were found to demonstrate seasonality depending on the country - namely 'hay fever', 'allergy' and 'pollen' - showing cultural differences. A single set of terms cannot be used across all European countries, but allergy seasonality can be compared across Europe providing the above three terms are used. Using longitudinal data in different countries and multiple terms, we identified an awareness-related spike of searches (December 2016).

terms were analysed: 'allergy' as a disease, 'hay fever' as a topic, 'asthma' as a disease, 'conjunctivitis' as a disease and 'pollen' as topic. "Rhinitis" as a term or 'allergic rhinitis' as a disease were labelled 'hay fever' as a topic. Only 'diseases' and 'topics' were automatically translated by GT, whereas 'terms' were not translated. In this study, we did not control for the translations of terms in different languages as we wanted a real-life assessment of GT. 'Rhinitis' as a term was not translated manually as certain countries are multilingual (e.g. Belgium and Switzerland).

Google Trends is based on Google Search. It shows how often a particular search term is entered relative to the total search volume across various regions of the world, and in various languages. The horizontal axis of the main graph represents time (starting from 2004), and the vertical axis shows how often a term is searched relative to the total number of searches globally.

We examined GT queries from 1 January 2011 to 20 December 2016 (using the standard 5-year analytic window of GT) for all 28 EU countries, Norway and Switzerland. We established visually (JB) patterns of countries according to the seasonality of terms (Table 1). We then compared the country patterns (based on seasons) with the mean 5-year terms provided by GT.

We did not compare the results of the 5-year trend with those of one year or less, as the goal of the study was not to define whether GT can be used as sentinel, but to assess whether there are differences between terms in European countries.

Results

Using the GT 5-year graph, an annual and clear seasonality of queries was found in all countries apart from Cyprus, Estonia, Latvia, Lithuania and Malta (Fig. S2). In these countries, some trends in seasonality were observed but they were not as clear as in the other countries.

Five patterns of country patterns were identified (Table 1). In 16 countries, a spike of queries was observed in the last week of December 2016. Figure 1 shows one example of each cluster, whereas all countries are presented in Fig. S1.

Country patterns were defined according to seasonality of terms. However, this classification fits with the mean levels of

Abbreviations

ARIA, Allergic Rhinitis and its Impact on Asthma; EU, European Union; GT, Google Trends; MACVIA, *Contre les MAIadies Chroniques pour un Vieillessement Actif*; MASK, MACVIA-ARIA Sentinel NetworkK.

Table 1 Patterns of countries

Cluster	Characteristics	Countries	Average means over 5 years			
			Allergy	Hay fever	Pollen	Asthma
A	Peak of 'hay fever' higher than 'allergy'	Ireland	45	12	13	24
		Netherlands	21	9	2	7
		UK	36	14	12	21
B	Peak of 'hay fever' lower than 'allergy' and higher than 'asthma'	Austria	38	5	4	9
		Belgium	49	6	7	10
		Denmark	39	3	7	14
		Germany	53	11	5	16
		Luxembourg	28	8	6	8
		Switzerland	51	11	10	14
C	Peak of 'pollen' only and higher than 'asthma'	Finland	53	1	8	22
		France	51	3	11	11
		Norway	51	1	7	16
		Sweden	43	1	8	13
D	Peak of 'hay fever' and 'pollen' lower than or similar to 'asthma' OR	Bulgaria	44	3	2	14
		Croatia	32	1	1	6
	Peak of 'allergy' without peak of 'hay fever' or 'pollen'	Czech Republic	40	2	2	10
		Greece	50	2	3	16
		Hungary	34	1	3	6
		Italy	46	2	1	7
		Poland	49	1	1	10
		Portugal	48	2	1	10
		Romania	51	1	1	10
		Slovenia	45	1	2	10
		Slovakia	40	4	3	9
		Spain	38	1	1	7
E	No season identified	Cyprus	39	8	8	17
		Estonia	54	3	5	15
		Latvia	58	2	5	16
		Lithuania	49	6	6	10
		Malta	44	15	13	26

queries over 5 years. In Cluster A countries, 'allergy' queries were reported at a lower level than 'hay fever' and did not show any clear seasonal pattern. In pattern B, C and D countries, 'allergy' queries were higher than the other terms and showed a seasonal pattern. When all countries with a seasonal pattern were examined, the peak of queries was the same for 'allergy' and 'hay fever' (Cluster B) or 'allergy' and 'pollen' (Cluster C). Moreover, in patterns A, B, C and D, the seasonal patterns were similar between countries for the five years (Figs 1 and S1).

Asthma and conjunctivitis did not show any seasonal pattern using the 5-year approach (data not shown for conjunctivitis).

There was a spike of queries in Austria, Belgium, Bulgaria, Croatia, Czech Republic, Estonia, Finland, Hungary, Ireland, Italy, Malta, Norway, Portugal, Romania, Spain and Switzerland at the end of December 2016 (Figs 1, S1 and Table S1). These queries appeared within a week in all of the countries and were unrelated to the seasons in previous years. The shape of the spike was steeper than during allergen seasons. Moreover, two new patterns were found (F: peak of 'allergy' and 'asthma' and G: peak of 'asthma'). The change in country patterns occurred in 10 (66.7%) of the countries.

Discussion

Google Trends reflects the proportion of specific queries in relation to all queries. Although terms were translated into the different languages using the automatic Google translator, the reporting of GT differs widely between countries. There are country-specific trends towards a different awareness of allergy terms. Different terms account for seasonality ('allergic rhinitis' or 'hay fever' as a topic, 'allergy' as a disease, 'pollen' as a topic). A single set of terms cannot be used across all countries, but allergy seasonality can be compared across Europe providing the three terms are used. Moreover, longitudinal data in different countries and multiple terms enabled a spike of terms to be identified, apparently unrelated to allergen exposure. The study in Europe is generalizable to other countries, and the same clustering approach may be used.

Google Trends does not provide a uniform, unbiased geographical sample. The data pool is skewed towards high-income countries with a higher access to computers and Internet. These problems are unlikely to be of importance in the European countries we studied. Moreover, GT does not allow a direct access to absolute numbers.




Figure 1 Examples of patterns. Cluster A: Ireland. Cluster B: Belgium. Cluster C: Finland. Cluster D: Croatia. Cluster E: Malta.

Significant spikes in the Google search were found with increased awareness of a disease. Examples include the death of a celebrity (13) and the heart transplantation in the month when Vice President Cheney underwent the respective procedure (14). It is likely that the December 2016 spikes were associated with the awareness of thunderstorm asthma (15). Two severe episodes occurred in Australia and Kuwait in the past month and were largely publicized in the media. However, using multiple terms, a 5-year follow-up and multiple countries, we were able to identify the awareness-related spikes.

The clustering approach based on seasonality was apparently sound as the mean levels of terms over the 5 years differed between patterns according to the seasonal patterns. Moreover, the patterns group countries as expected: Ireland and the UK in Cluster A (with the Netherlands), German-speaking countries in Cluster B and three Nordic countries in Cluster C. These patterns may be related to the understanding of the terms in different languages or to cultural differences. However, combining the three terms, we could assess seasonal changes in most countries. It is interesting to note that the seasons occur at a similar time (with some small differences due to climate) across Europe, in particular for the birch, grass and ragweed pollen seasons. The clustering approach has improved our knowledge on the apparent awareness spikes of December 2016.

In this study, we did not assess antihistamine queries or the different pollen species (e.g. birch, grass, *Parietaria* or ragweed) as the aim was to assess terminology. Moreover, we did not study yearly queries which may indicate some seasonality for asthma and a more precise seasonality in the five countries of Cluster E.

This study has several potential impacts (Table 2).

The results of this study will be used in the development of the MASK sentinel network (1). MASK (MACVIA (*Contre les Maladies Chroniques pour un Vieillissement Actif*)-ARIA Sentinel Network) uses mobile technology to develop care pathways for the management of rhinitis and asthma by a multidisciplinary group or by patients themselves (16). An app (Android and iOS) is available in 20 countries and 15

Table 2 Potential impact of the study

- It is possible to investigate Google search trends across a range of European countries
- Although patients in Europe overall experience the same symptoms of allergic rhinitis induced by pollens which induce an allergic reaction, they use searches which differ between countries. We have found that the searches are similar in different regions of the same country. The clusters we have proposed are in line with cultural differences.
- When GTs are used in allergic rhinitis, patients in most European countries use other search terms.
- There is a need to use multiple searches for a single disease.
- The results call for uniform nomenclature and self-management guidelines to improve detection and proactive treatment of seasonal allergy symptoms.
- GT analysis is a tool to improve and tailor our communication to the patients and general public besides building our sentinel network

languages. It uses a visual analogue scale to assess symptom control and work productivity as well as a clinical decision support system. The scaling up strategy uses the recommendations of the European Innovation Partnership on Active and Healthy Ageing (17).

More data are required to investigate the importance of GT in a sentinel network, but it is clear that a combination of three terms over a period of observation is needed across Europe to establish baseline and to assess spikes.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1 Spike of queries at the end of December 2016.

Figure S1 Google Trends for the past 5 years in Europe.

References

1. Bousquet J, Schunemann HJ, Fonseca J, Samolinski B, Bachert C, Canonica GW et al. MACVIA-ARIA Sentinel Network for allergic rhinitis (MASK-rhinitis): the new generation guideline implementation. *Allergy* 2015;**70**:1372–1392.
2. Pfaar O, Bastl K, Berger U, Buters J, Calderon MA, Clot B et al. Defining pollen exposure times for clinical trials of allergen immunotherapy for pollen-induced rhinoconjunctivitis – an EAACI Position Paper. *Allergy* 2017;**72**:713–722.
3. Bastl K, Kmenta M, Pessi AM, Prank M, Saarto A, Sofiev M et al. First comparison of symptom data with allergen content (Bet v 1 and Phl p 5 measurements) and pollen data from four European regions during 2009–2011. *Sci Total Environ* 2016;**548–549**:229–235.
4. Buters JT, Weichenmeier I, Ochs S, Pusch G, Kreyling W, Boere AJ et al. The allergen Bet v 1 in fractions of ambient air deviates from birch pollen counts. *Allergy* 2010;**65**:850–858.
5. Myszkowska D, Majewska R. Pollen grains as allergenic environmental factors—new approach to the forecasting of the pollen concentration during the season. *Ann Agric Environ Med* 2014;**21**:681–688.
6. de Weger LA, Beerthuis T, Hiemstra PS, Sont JK. Development and validation of a 5-day-ahead hay fever forecast for patients with grass-pollen-induced allergic rhinitis. *Int J Biometeorol* 2014;**58**:1047–1055.
7. König V, Mosges R. A model for the determination of pollen count using google search queries for patients suffering from allergic rhinitis. *J Allergy (Cairo)* 2014;**2014**:381983.
8. Willson TJ, Lospinoso J, Weitzel E, McMains K. Correlating regional aeroallergen effects on internet search activity. *Otolaryngol Head Neck Surg* 2015;**152**:228–232.
9. Zuckerman O, Luster SH, Bielory L. Internet searches and allergy: temporal variation in regional pollen counts correlates with Google searches for pollen allergy related

- terms. *Ann Allergy Asthma Immunol* 2014;**113**:486–488.
10. Willson TJ, Shams A, Lospinoso J, Weitzel E, McMains K. Searching for cedar: geographic variation in single aeroallergen shows dose response in internet search activity. *Otolaryngol Head Neck Surg* 2015;**153**:770–774.
 11. Kang MG, Song WJ, Choi S, Kim H, Ha H, Kim SH et al. Google unveils a glimpse of allergic rhinitis in the real world. *Allergy* 2015;**70**:124–128.
 12. Bousquet J, Hellings PW, Agache I, Bedbrook A, Bachert C, Bergmann KC et al. ARIA 2016: care pathways implementing emerging technologies for predictive medicine in rhinitis and asthma across the life cycle. *Clin Transl Allergy* 2016;**6**:47.
 13. Bragazzi NL, Watad A, Brigo F, Adawi M, Amital H, Shoenfeld Y. Public health awareness of autoimmune diseases after the death of a celebrity. *Clin Rheumatol*. 2016; doi: 10.1007/s10067-016-3513-5
 14. Pandey A, Abdullah K, Drazner MH. Impact of Vice President Cheney on public interest in left ventricular assist devices and heart transplantation. *Am J Cardiol* 2014;**113**:1529–1531.
 15. Thunderstorm asthma. en.wikipedia.org/wiki/Thunderstorm_asthma. 2016.
 16. Bousquet J, Addis A, Adcock I, Agache I, Agusti A, Alonso A et al. Integrated care pathways for airway diseases (AIRWAYS-ICPs). *Eur Respir J* 2014;**44**:304–323.
 17. Bousquet J, Farrell J, Crooks G, Hellings P, Bel EH, Bewick M et al. Scaling up strategies of the chronic respiratory disease programme of the European Innovation Partnership on Active and Healthy Ageing (Action Plan B3: Area 5). *Clin Transl Allergy* 2016;**6**:29.